

# Inform all eligible patients of the vaccines they need



As a trusted provider, you can help  
close the vaccination gap



# Assess patient's eligibility for vaccines

Evaluate based on patient's age, risk factors, and vaccination history

Download CDC Vaccine Schedules app for more information		Vaccine				
		RSV <sup>1*</sup>	Pneumococcal <sup>††</sup>	2024-2025 COVID-19 <sup>††</sup> Additional doses may be recommended	Shingles <sup>§§</sup>	Tdap <sup>¶¶</sup>
Age	19-49 <sup>#</sup>			●		●
	19-49, risk factors <sup>#  **</sup>		●	●	●	●
	50-59 <sup>#</sup>		●	●	●	●
	60-74 <sup>#</sup>		●	●	●	●
	60-74, risk factors <sup>#  </sup>	●	●	●	●	●
	75+ <sup>#</sup>	●	●	●	●	●

Complete the tear pad with recommended vaccines, using it as a counseling tool and, if needed, an appointment reminder for follow-up vaccinations 

\*CDC recommends everyone 75+ and people 60-74 with certain medical conditions and/or risk factors receive a single dose of RSV vaccine. Risk factors may include heart failure, asthma, diabetes, or immunocompromising conditions. Download the **CDC Vaccine Schedules** app for more information on risk factors.<sup>1</sup>

†CDC recommends pneumococcal vaccination for adults 50+ or 19+ with certain medical conditions and/or risk factors. Download the **CDC Vaccine Schedules** app for more information on risk factors and the number of doses recommended based on vaccination history.<sup>1</sup>

‡CDC recommends adults get a 2024-2025 COVID-19 vaccine, including people who previously received a COVID-19 vaccine or had COVID-19. For adults 65+ and individuals who are moderately or severely immunocompromised, download the **CDC Vaccine Schedules** app for more information on additional doses.<sup>1</sup>

§CDC recommends everyone 50+ and those with immunocompromising conditions get shingles vaccine as a 2-dose series, administered 2-6 months apart.<sup>1</sup>

¶CDC recommends adults get a Tdap vaccine every 10 years.<sup>1</sup>

#Risk factors for Tdap may include pregnancy and wounds. Download the **CDC Vaccine Schedules** app for more information.<sup>1</sup>

||Risk factors for RSV and pneumococcal may include heart failure, asthma, diabetes, and immunocompromising conditions. Download the **CDC Vaccine Schedules** app for more information on vaccine-specific risk factors.<sup>1</sup>

\*\*Risk factors for shingles may include transplant, cancer, HIV, and autoimmune & inflammatory conditions.<sup>2</sup>



# Vaccine Action Plan

Save time by getting more than 1 vaccine in the same visit. Talk to your pharmacist.

Name: \_\_\_\_\_ Age: \_\_\_\_\_

Chronic condition(s):\*  Yes  No

Vaccine(s):  RSV  Pneumococcal  COVID-19  Shingles  Tdap

Scheduled Visit Date: \_\_\_\_\_

Time: \_\_\_\_\_

Pharmacy Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Help protect yourself by making a plan to  
receive all recommended vaccines



COVID-19 = coronavirus disease 2019; RSV = respiratory syncytial virus; Tdap = tetanus, diphtheria, and pertussis.

\* Ask your pharmacist about how risk factors such as heart failure, asthma, diabetes, or a weakened immune system may affect your vaccination plan.



Empower patients to take control of their well-being and get vaccinated today



CDC = Centers for Disease Control and Prevention; COVID-19 = coronavirus disease 2019; HIV = human immunodeficiency virus; RSV = respiratory syncytial virus; Tdap = tetanus, diphtheria, and pertussis.

**References:** 1. Centers for Disease Control and Prevention (CDC). Recommended adult immunization schedule. Updated November 21, 2024. Accessed January 6, 2025. <https://www.cdc.gov/vaccines/hcp/imz-schedules/downloads/adult/adult-combined-schedule.pdf> 2. CDC. Clinical considerations for Shingrix use in immunocompromised adults aged  $\geq 19$  years. Updated July 9, 2024. Accessed January 24, 2025. <https://www.cdc.gov/shingles/hcp/vaccine-considerations/immunocompromised-adults.html>